
Cosmic Frontier Computing Collaboration

ASCR SciDAC Institutes


Collaboration Codes

Cosmological Surveys


 Mysterious substance contributing most of the energy in the Universe




- Mysterious substance contributing most of the energy in the Universe
- Energy density remains constant as Universe expands!




- Mysterious substance contributing most of the energy in the Universe
- Energy density remains constant as Universe expands!
- Energy associated with empty space? [Recall Heisenberg]




- Mysterious substance contributing most of the energy in the Universe
- Energy density remains constant as Universe expands!
- Energy associated with empty space? [Recall Heisenberg]
- Answer wrong by 120 orders of magnitude



- Mysterious substance contributing most of the energy in the Universe
- Energy density remains constant as Universe expands!
- Energy associated with empty space? [Recall Heisenberg]
- Answer wrong by 120 orders of magnitude
- Need to study observationally



Gravitational Lensing



- Light paths distorted by intervening structure
- Distortions depend on both distance-redshift relation and growth of structure
- Both of these things are affected by the nature of dark energy

Galaxy Ellipticities

Spherical galaxy

Mass

Galaxy Ellipticities

Spherical galaxy





Mass

Galaxy Ellipticities

Spherical galaxy



Mass




Galaxy appears elliptical




Extracting information from galaxy ellipticities

- Mean ellipticity is zero
- Information is contained in variance and correlations



Quantification: Spectra






Hoekstra H, Jain B. 2008.

Annu. Rev. Nucl. Part. Sci. 58:99-123

These spectra computed neglecting baryons ("N-Body")






Hoekstra H, Jain B. 2008.


Annu. Rev. Nucl. Part. Sci. 58:99-123

Need to account for baryons ("Hydro")



Baryonic Effects


- Differences large: red
 & blue curves outside
 projects errors from
 upcoming surveys
- Effect uncertain:
 different
 implementations give
 vastly different results



Rudd et al. 2007

Dark Energy Survey

Identified as one of key systematics by DES Weak Lensing Working Group



Study with latest Simulations (Zentner, Semboloni, Dodelson, Eifler, Kraus, Hearin)


	Box Size (Mpc)	N	Mass resol	Spatial resol
Jing et al 2006	100	512	7e8	?
Rudd, Zentner, Kravtsov 2008	60	256	1e9	2 kpc
Guilett, Teyssier, Colombi 2010	50	1024	1e7	1 kpc
van Daalen et al 2011 (OWLS)	100	512	7e8	2 kpc

OverWhelmingly Large Simulations (OWLS)


~10 simulations with same initial conditions/resolution/box size, but different prescriptions for baryons

Simulation	Description
AGN	Includes AGN (in addition to SN feedback)
AGN_WMAP7	Same as AGN , but with a WMAP7 cosmology
DBLIMFV1618	Top-heavy IMF at high pressure, extra SN energy in wind velocity
DMONLY	No baryons, cold dark matter only
$DMONLY_WMAP7$	Same as DMONLY, but with a WMAP7 cosmology
MILL	Millennium simulation cosmology (i.e. WMAP1), $\eta=4$ (twice the SN energy of REF)
NOSN	No SN energy feedback
$NOSN_NOZCOOL$	No SN energy feedback and cooling assumes primordial abundances
NOZCOOL	Cooling assumes primordial abundances
WDENS	Wind mass loading and velocity depend on gas density (SN energy as REF)
WML1V848	Wind mass loading $\eta = 1$, velocity $v_{\rm W} = 848 {\rm km s^{-1}}$ (SN energy as REF)
WML4	Wind mass loading $\eta=4$ (twice the SN energy of REF)

Baryonic Effects at least as large as in Rudd et al.




How bad is ignoring the effect? (Raw Bias)




Parameter 1

Raw Bias is significantly larger than statistical error for many of the simulations




How can we mitigate this?


Recall from Nick's talk that baryons tend to make halos more concentrated




Mitigation Strategy

- Introduce new parameters (beyond the standard cosmological parameters) that model the new massconcentration relation
- Fit for all parameters (cosmo + Mass-concentration)
- Errors will go up (green larger than blue) but residual bias should go down






Parameter 1



Residual Bias smaller than Statistical error even if small angular scales are included



Exciting Plans

	Box Size (Mpc)	N	Mass resol	Spatial resol
Jing et al 2006	100	512	7e8	?
Rudd, Zentner, Kravtsov 2008	60	256	1e9	2 kpc
Guilett, Teyssier, Colombi 2010	50	1024	1e7	1 kpc
van Daalen et al 2011 (OWLS)	100	512	7e8	2 kpc
Our plan	200	1024+	7e8	<1 kpc

We [Nick Gnedin, Scott Dodelson, John Freeman, Qiming Lu] look forward to working with you!